

**Summary of marine mammal
observations during 2011 surveys**

Massachusetts Water Resources Authority
Environmental Quality Department
Report 2013-03



Citation

Wu D. 2013. **Summary of marine mammal observations during 2011 surveys.** Boston: Massachusetts Water Resources Authority. Report 2013-03. 14 pp.

SUMMARY OF MARINE MAMMAL OBSERVATIONS DURING 2011 SURVEYS

**Massachusetts Water Resources Authority
Charlestown Navy Yard
100 First Avenue
Boston, MA 02129**

Prepared by
David Wu

Technical Report No: 2013-03

Acknowledgements

Marine mammal observers were individually contracted to assist Battelle in collecting the data contained in this report. The dedication and professionalism of David Silvia is appreciated.

Thanks and appreciation are also extended to the captains, crews, and scientific personnel of all the research vessels involved with the surveys of Boston Harbor and Massachusetts Bay.

Thanks to Matt Fitzpatrick of Battelle for compiling the mammal data.

TABLE OF CONTENTS

1.0	Introduction.....	1
2.0	Background.....	1
3.0	Methods.....	4
4.0	Results.....	8
5.0	Discussion.....	11
6.0	References.....	12

LIST OF TABLES

Table 1.	Year 2011 effluent outfall ambient monitoring surveys and <i>Alexandrium</i> surveys when marine mammals were sighted.	8
Table 2.	Year 2011 Boston Harbor and Massachusetts Bay shellfish water quality monitoring surveys when marine mammals were sighted.....	9

LIST OF FIGURES

Figure 1.	MWRA effluent outfall water column monitoring stations.....	5
Figure 2.	MWRA <i>Alexandrium</i> monitoring stations.....	6
Figure 3.	MWRA Boston Harbor and shellfish-growing water quality monitoring stations.....	7
Figure 4.	Locations of whale sightings during 2011 surveys.....	10

1.0 Introduction

At least five endangered species of whales are known to visit or inhabit the Massachusetts and Cape Cod Bay area (Environmental Protection Agency [EPA] 1993): the right whale, humpback whale, finback whale, sei whale (rarely observed) and blue whale (rarely observed). Several non-endangered marine mammal species are also found: minke whales, pilot whales, harbor porpoises, Atlantic white-sided dolphins, white beaked dolphins, hooded seals, harp seals, gray seals, and harbor seals.

Since 1995, Massachusetts Water Resources Authority (MWRA) has included marine mammal observers on monitoring surveys. The MWRA surveys are being conducted as part of the long-term Harbor and Outfall Monitoring Project designed to verify compliance with the discharge permit and to assess the potential environmental impact of treated sewage effluent discharge into Massachusetts Bay. These observers were included in response to a National Marine Fisheries Service request that MWRA provide observational data and set a positive example by using observers to minimize the chances of collision with a right whale. In addition to looking for right whales, observers noted other marine mammals. On surveys where observers were not present, the chief scientist and field crew documented any incidental sightings of marine mammals.

Marine mammal observers were present on all effluent outfall water quality surveys during 2011. Observers were not present on Boston Harbor surveys, floatable debris surveys, and bacteria surveys for shellfish water quality monitoring.

2.0 Background

A brief description of when marine mammals are expected to be found in Massachusetts and Cape Cod Bays is presented and discussed below.

The right whale (*Eubalaena glacialis*) is critically endangered. Based on historical sightings, right whales can be expected to visit Massachusetts and Cape Cod Bays throughout the year (Brown *et al.* 2002), with peak abundance in February, March and early April (Hamilton and Mayo 1990). Approximately 70% of the catalogued population of right whales have been reported to visit Cape Cod Bay and Massachusetts Bay (Brown *et al.* 2002), and NMFS has designated the Bays as an “area of high use” (NMFS 2013a). The use of the eastern portion of Stellwagen Bank/Wildcat Knoll by right whales has been noted during extended surveys by the Center for Coastal Studies (Brown *et al.* 2002). The total population of right whales in the western Atlantic Ocean is estimated to be about 400 individuals (NMFS 2013a).

The humpback whale (*Megaptera novaeangliae*) is an endangered species of whale known to feed within the Gulf of Maine in the spring, summer and fall (Waring *et al.* 1999). In the winter, some, but not all, humpbacks will migrate to mating and calving grounds in the West Indies (NMFS 2013a). Historic records indicate that humpbacks have been documented on Stellwagen Bank from April through December (CeTap 1982; Geraci *et al.* 1989; NMFS 1991). However, distribution appears to correlate with prey densities (Waring *et al.* 1999). The amount of humpback whale use of the Stellwagen area varies periodically most likely based on the availability of sand lance as prey (Payne *et al.* 1986; Payne *et al.* 1990; Weinrich *et al.* 1997). The best available estimate of the humpback population in the western Atlantic is about 11,500 individuals (NMFS 2013a).

The finback (or fin) whale (*Balaenoptera physalus*) is considered to be an endangered species and is the most abundant and frequently sighted of the endangered whales that visit Massachusetts and Cape Cod Bays (EPA 1993). Finbacks are sighted year round in the Stellwagen Bank area with a peak abundance occurring between the spring and fall (Pett and McKay 1990). Finbacks do also migrate, potentially from

the North Atlantic to the West Indies, but migratory routes are unknown. Insufficient data exists to determine population trends of the western North Atlantic population, but the minimum number of individuals is estimated at 1,678 whales (NMFS 2013a).

The sei whale (*Balaenoptera borealis*) and blue whale (*Balaenoptera musculus*) are both endangered species (EPA 1993). The sei whale is uncommon but is regularly sighted (Schilling *et al.* 1992), while the blue whale is rarely sighted in Massachusetts and Cape Cod Bays (EPA 1993). For blue whales, Massachusetts and Cape Cod Bays may represent the southern limit of their feeding area (NMFS 2013a). Both blue and sei whales typically remain in deeper water (more than 100 meters) and further offshore (CeTap 1982). However, sightings of these species in coastal areas may correspond to changes in prey distribution (Payne *et al.* 1990, Wenzel *et al.* 1988). In general, though, the large-scale distribution and movement patterns of sei whales are not well known. There are no current population estimates for sei whales, although the North Atlantic population seems to be “relatively abundant” (NMFS 2013) after intensive whaling in the 19th and 20th centuries. An estimated 400-600 blue whales reside in all of the North Atlantic (NMFS 2013a).

The minke whale (*Balaenoptera acutorostrata*) is a non-endangered species typically seen in the Stellwagen Bank area during the spring, summer and fall (CeTap 1982; Pett and McKay 1990). During the winter, minke whale sightings in New England appear to decline dramatically (Waring *et al.* 1999). For management purposes, New England minke whales are known as the Canadian Eastern Coastal Stock, with an estimated 2,500-3,000 individuals. Insufficient data exists for analysis of population trends (NMFS 2013a).

The Atlantic white-sided dolphin (*Lagenorhynchus acutus*) is a species of dolphin found from central west Greenland to North Carolina (Waring *et al.* 1999). The Gulf of Maine stock of Atlantic white-sided dolphins is classified as strategic by the National Marine Fisheries Service (Waring *et al.* 1999). Sightings of these dolphins in the Stellwagen Bank and Cape Cod Bay areas are common in the spring, summer (Weinrich *et al.* 2001), and, to a lesser extent, the fall (Pett and McKay 1990). The western Atlantic population is estimated at 63,000 individuals, with insufficient information to determine population trends (NMFS 2013a).

The Atlantic pilot whale or long-finned pilot whale (*Globicephala melas*) is the largest species of dolphin found in cool temperate waters off Labrador, Newfoundland, and in the St. Lawrence River with sporadic sightings as far south as Maryland and Virginia (Bulloch 1993). Pilot whales form schools of a few to many hundreds of individuals and are mainly found relatively close to shore. Pilot whale distribution and abundance appear to be linked to the topography of the sea floor and the abundance of their primary food source, squid (Harrison and Bryden 1989). Population estimates of the long-finned pilot whale in the western North Atlantic range from 25,000 to 31,000 individuals. Note that this estimate includes short-finned pilot whales (*Globicephala macrorhynchus*), a different species that is almost visually indistinguishable from long-finned pilot whales in the field (NMFS 2013a).

The gray seal (*Halichoerus grypus*) is a non-endangered species of pinniped found from Maine to Long Island Sound (Rough 1995). A small, year round breeding population is known to occur on outer Cape Cod and Nantucket Island (Waring *et al.* 1999). The majority of gray seal sightings in Cape Cod Bay and the Stellwagen Bank area occur during the winter and spring, although periodic sightings have been recorded in the summer (Center for Coastal studies unpublished data). Current population in the western North Atlantic is estimated at about 250,000 individuals. Most recent surveys seem to indicate that population is increasing after a long period of decline due to hunting for both subsistence and fur (NMFS 2013a).

Harbor porpoises (*Phocoena phocoena*) of the Gulf of Maine/Bay of Fundy stock are classified as strategic by the National Marine Fisheries Service (Waring *et al.* 1999). Historic data indicate that harbor porpoises can be found in the Stellwagen Bank area and Cape Cod Bay from December through June (Pett and McKay 1990). A 2006 aerial survey of the Gulf of Maine/Bay of Fundy stock estimated 89,054 individuals. No population trend analysis has been performed (NMFS 2012).

The harbor seal (*Phoca vitulina*) is a non-endangered species of pinniped commonly found in the near shore waters around New England (Katona *et al.* 1993). Harbor seals are most frequently seen in the Stellwagen Bank and Cape Cod Bay areas in the winter and early spring with sightings beginning in late September (Pett and McKay 1990). No surveys of population have been conducted since 2001 but between the passage of the Marine Mammal Protection Act in 1972 and the 2001 survey populations in New England increased (NMFS 2013b).

3.0 Methods

Figure 1 shows MWRA effluent outfall ambient monitoring water column sampling stations. The year 2011 was the first year the second revision of the ambient monitoring plan design was implemented (MWRA 2010). The revised design focuses more on stations likely to be impacted by the outfall; there are fewer distant reference stations. The total number of outfall monitoring stations was reduced from 33 to 14. Also, the number of surveys was changed from 12 nearfield and six farfield surveys annually to nine surveys annually of all 14 stations. These changes have meant that the surveys were generally completed in a single day while previously multiple days were needed to accomplish the farfield surveys. Thus, there was less time to observe marine mammals than in previous years. In addition, except for the *Alexandrium* surveys, MWRA's marine mammal observations no longer include the areas where whales are most frequently found (Stellwagen Bank National Marine Sanctuary and Cape Cod Bay). Those areas are monitored specifically for marine mammals under separate non-MWRA research programs.

Marine mammal observations were performed during all daylight hours while transiting between stations during water column surveys, and while the vessel was on-station for sampling operations. During vessel transits, the observer continuously scanned the sea surface from directly ahead to 90 degrees abeam on either side of the vessel. Initial sightings were made by eye with confirmation and identification aided by binoculars. While on-station, the observer scanned 360 degrees around the vessel. The observer was typically positioned at the highest secure vantage point of the survey vessel. Weather conditions, safety of the observer, and limiting interference with the operation of the vessel and sampling team were all factors that influenced the position of the observer on board the vessel.

Two survey vessels were used as observation platforms during the course of the year. The observer's eye-height above the sea surface was approximately 4 meters on the R/V *Tioga* and R/V *Andy Lynn VI* and 2.5 meters aboard the R/V *Aquamonitor*. Observations were conducted 40 minutes out of every hour and were suspended when visibility was reduced to zero or when darkness occurred. The vessels R/V *Auk* and R/V *Tioga* were also used for surveys with onboard marine mammal observers.

For some surveys, dedicated marine mammal observers were not present. The scientific crew on board the R/V *Merganser* and R/V *Aquamonitor* observed marine mammals while on these surveys. These vessels were used to conduct MWRA Boston Harbor surveys and some other nearfield, farfield, and *Alexandrium* rapid response surveys (Figures 2 and 3). Similar to previous years, data from those surveys are included in this report.

Vessel track, station sequence, and number of stations varied among cruises, due to the constraints of weather, special survey requirements, or both.




Figure 1. MWRA effluent outfall water column monitoring stations.




Figure 2. MWRA *Alexandrium* monitoring stations




Figure 3. MWRA Boston Harbor and shellfish-growing water quality monitoring stations

4.0 Results

Observation of marine mammals on surveys designed and operated for the collection of water quality data places limitations and constraints on the method of observation and on the conclusions that may be drawn from the data. Standard line transect methodology is not possible on such surveys, and different vessels (which vary the characteristics of the survey platform) were used during the year. Therefore, it is not appropriate to use these opportunistic sightings to estimate animal abundance. The data provide useful qualitative information concerning seasonal patterns and relative abundance within the same study area.

During the 2011 monitoring year, there were nine effluent outfall ambient monitoring surveys (WN) and three *Alexandrium* surveys (AF), 160 Boston Harbor water quality surveys, and twelve Massachusetts Bay shellfish water quality monitoring surveys. Survey team members counted six individual whales including five minke whales and one finback whale. Also counted were approximately 50 short beaked common dolphins, 12 harbor porpoises, 30 harbor seals, and one unidentified seal. Tables 1 and 2 summarize the locations and dates of all MWRA's sightings of whales, pinnipeds and dolphins in 2011. The locations of whale sightings are shown in Figure 4.

Table 1. Year 2011 effluent outfall ambient monitoring surveys and *Alexandrium* surveys when marine mammals were sighted.

Survey ID	Date/Time	Number	Mammal	Location	Sighting Comments	Observer Present
WN112 <i>R/V Aquamonitor</i>	3/15/11 0805	1	Harbor seal	42.3383, -70.942	Vicinity of F23	Yes
WN113 <i>R/V Aquamonitor</i>	4/7/11 0720	10	Harbor seal	42.27163, -70.93028	Near Grape Island	Yes
	4/7/11 1108	1	Minke whale	42.3679, -70.77863	Between N21 and N18	Yes
AF111 <i>R/V Aquamonitor</i>	5/25/11 1523	1	Harbor porpoise	42.1254, -70.54577	In transit from F06 to AF2	No
	5/25/11 1523	1	Harbor porpoise	42.0543, -70.544	In transit from AF2 to AF1	No
AF113 <i>R/V Aquamonitor</i>	6/8/11 0830	1	Seal	42.39223, -70.754916	At N16	No
	6/8/11 1142	1	Minke whale	42.44443, -70.51318	In transit from F22 to AF6	No
WN115 <i>R/V Aquamonitor</i>	6/16/11 1610	1	Harbor seal	42.325, -70.862	Between N10 and Nantasket Roads entrance	Yes
WN116 <i>R/V Aquamonitor</i>	7/19/11 0910	1	Minke whale	42.4795, -70.617	Vicinity of F22	Yes
WN117 <i>R/V Aquamonitor</i>	8/18/11 1340	1	Minke whale	42.17183, -70.5783	Near F06	Yes
WN119 <i>R/V Aquamonitor</i>	10/18/11 0830	3	Harbor seal	42.314, -70.908	On Toddy Rocks, Hull	Yes

Table 2. Year 2011 Boston Harbor and Massachusetts Bay shellfish water quality monitoring surveys when marine mammals were sighted.

Survey ID	Date/Time	Number	Mammal	Location	Sighting Comments	Observer Present
WQM2011 <i>R/V Merganser</i>	1/5/11 0845	1	Harbor seal	42.375159, -71.047275	Inner Harbor, near Pier 8	No
PC112 <i>R/V Merganser</i>	2/7/11 0945	1	Harbor seal	42.2683333, -70.73500	Near F13	No
WQM2011 <i>R/V Merganser</i>	3/23/11 1114	1	Harbor seal	42.294070, -70.91793	Between Peddock's Island and Hull	No
	3/23/11 1125	1	Harbor seal	42.31794, -70.93180	Just W of George's Island	No
WQM2011 <i>R/V Merganser</i>	4/7/11 0817	1	Harbor seal	42.38516, -71.05145	Upstream of Tobin Bridge	No
	4/7/11 0850	2	Porpoise	42.38271, -71.04678	Under Tobin Bridge	No
	4/7/11 1010	1	Harbor porpoise	42.33689, -71.00478	Near 038	No
WQM2011 <i>R/V Merganser</i>	4/19/11 1228	1	Harbor seal	42.33253, -70.95903	Off 106	No
PC114 <i>R/V Merganser</i>	4/25/11 1000	1	Harbor seal	42.37687, -70.73290	Between N07 and N16	No
	4/25/11 1118	1	Harbor porpoise	42.43347, -70.84969	Between N02 and F18	No
	4/25/11 1120	5	Harbor porpoises	42.43480, -70.85487	Between N02 and F18	No
PC115 <i>R/V Merganser</i>	5/2/11 1127	2	Harbor porpoises	42.4421667, -70.888333	At F18	No
PC116 <i>R/V Merganser</i>	6/6/11 1100	1	Minke whale	42.4361, -70.74073	Between N16 and N04	No
PC118 <i>R/V Merganser</i>	8/2/11 1012	1	Finback whale	42.4438333, -70.7365	N04	No
WQM2011 <i>R/V Merganser</i>	8/17/11 0846	1	Harbor seal	42.37153, -71.04743	Inner Harbor, near Pier 7	No
PC11B <i>R/V Merganser</i>	11/8/11 0822	1	Harbor seal	42.30225, -70.80663	Near F14	No
	11/8/11 1126	1	Harbor seal	42.4421667, -70.888333	At F18	No
CSO2011 <i>R/V Merganser</i>	11/9/11 0808	1	Harbor seal	42.335, -70.9815	065	No
CSO2011 <i>R/V Merganser</i>	11/30/11 1103	1	Harbor seal	42.3859045, -71.0687348	069	No
PC11C <i>R/V Merganser</i>	12/5/11 1030	~50	Short beaked common dolphins	42.42204, -70.79391	Between N04 and N20	No
WQM2010 <i>R/V Merganser</i>	12/6/11 0853	1	Harbor seal	42.37536, -71.04542	Inner Harbor, near Pier 8	No
CSO2010 <i>R/V Merganser</i>	12/22/11 0930	1	Harbor seal	42.38467, -71.03166	Chelsea Creek	No




Figure 4. Locations of whale sightings during 2011 surveys

Note: The data displayed in this figure come from Tables 1 and 2 of this report.

5.0 Discussion

Unlike statistically-based programs or programs that are specifically designed to search for whales, the MWRA sightings are opportunistic and do not follow dedicated and systematic line transect methodology. Therefore, observations are descriptive and not a statistically robust population census. As noted above, the hours spent on the water were substantially reduced in 2011 compared to previous years, and the prime whale habitats of Stellwagen Bank and Cape Cod Bay are no longer included in MWRA's marine mammal observations.

MWRA's previous marine mammal reports compared the sightings of species of whales across areas surveyed and years (e.g., Wu 2011, Table 3). Although not identical, the best historical comparisons for 2011 whale observations would be with the past nearfield (NF) observations. From 1998-2010 the 13-year NF observations were: right whales total = 0; humpback whales total = 3, range 0-2/year; finback whales total = 8, range 0-3/year; minke whales total = 26, range 0-4/year; unidentified whales total = 14, range = 0-5/year. In 2011, MWRA observed 1 finback and 5 minke whales, which is in the range of previous NF observations.

Most observations of seals are when the vessels are transiting to and from the outfall monitoring area or during Boston Harbor surveys. The seals were typically resting upon rocks. During 2011, 31 pinnipeds were sighted. Thirty of these were harbor seals and one was a seal of unidentified species. These sightings were a decrease from 2010 when 90 pinnipeds were reported. For comparison, the numbers for 2001 to 2009 ranged from 76 to 303/year. Before 2001, 20 to 60 pinniped sightings were made throughout the survey area.

Approximately 50 short beaked common dolphins were sighted in 2011, in addition to 12 harbor porpoises.

MWRA no longer tabulates whale observations in Cape Cod Bay. Beginning in 2011, MWRA's Cape Cod Bay water quality monitoring is carried out by the Provincetown Center for Coastal Studies (PCCS), which has a long-standing scientific monitoring program for whales in Cape Cod Bay. Since 1998, PCCS has conducted systematic surveys of Cape Cod Bay and adjacent waters from January through mid-May. In 2010 PCCS (Stamieszkin *et al.* 2010) counted 163 different right whales identified using photographs. This number is comparable to sightings in 2007, 2008, and 2009. Half the individuals sighted in 2007 were seen again in the 2008 surveys, and 61% of the individuals seen in 2008 were spotted in 2009. From 2007 to 2010 at least 45% of the known right whale population has been sighted annually in the Cape Cod Bay, making it an important habitat for right whales (Stamieszkin *et al.* 2010, Leeney *et al.*, 2008, 2009).

6.0 References

Brown MW, O Nichols, MK Marx, and JN Ciano. 2002. Surveillance monitoring and management of North Atlantic right whales (*Eubalaena glacialis*) in Cape Cod Bay, Massachusetts: 2002. Final report to Division of Marine Fisheries, Commonwealth of Massachusetts, and Massachusetts Environmental Trust, September, 2002. 28 pp.

Bulloch DK. 1993. The Whale-Watcher's Handbook: A Field Guide to the Whales, Dolphins, and Porpoises of North America. Lyons & Burford, New York, NY. 114 pp.

CeTAP. 1982. A Characterization of Marine Mammals and Turtles in the Mid- and North Atlantic Areas of the U.S. Outer Continental Shelf. Final Report of the Cetacean and Turtle Assessment Program, University of Rhode Island, Kingston, RI. U.S. Dept. of the Interior, Bureau of Land Management, Washington, DC. Contract AA551-CT-48. 450 pp.

Environmental Protection Agency. 1993. Assessment of Potential Impact of the MWRA Outfall on Endangered Species. Boston, MA: U.S. Environmental Protection Agency.

Geraci, J.R., Anderson, D.M., Timperi, R.J., Early, G.A., Prescott, J.H., and Mayo, C.A. 1989. Humpback whales (*Megaptera novaeangliae*) fatally poisoned by dinoflagellate toxin. J. Fish. Res. Bd. Canada 46: 1895-1898.

Harrison R, MM Bryden. 1989. Whales, Dolphins and Porpoises. Weldon Owen Pty Limited, McMahons Point, Australia. 240 pp.

Jaquet, N., CA Mayo, D Osterberg, OC Nichols, MK Marx, and CL Browning. 2006. Surveillance, monitoring and management of North Atlantic right whales in Cape Cod Bay and adjacent waters – 2006. Final report submitted to the Division of Marine Fisheries, Commonwealth of Massachusetts. November 2006.

Jaquet N., Mayo C., Osterberg D., Browning C. L. and Marx M. K. 2007. Surveillance, Monitoring and Management of North Atlantic Right Whales in Cape Cod Bay and Adjacent Waters - 2007. Final report submitted to the Division of Marine Fisheries, Commonwealth of Massachusetts. October 2007:112pp, retrieved from <http://www.coastalstudies.org/what-we-do/right-whales/rwreports.htm>

Katona SK, V Rough, DT Richardson. 1993. A Field Guide to Whales, Porpoises, and Seals from Cape Cod to Newfoundland. Smithsonian Institution Press. Washington, DC. 316 pp.

Langton RW, JB Pierce, JA Gibson. 1994. Selected Living Resources, Habitat Conditions, and Human Perturbations of the Gulf of Maine. NOAA Tech. Memo NMFS-NE-106; 70 pp.

Leeney RH, K Stamieszkin, N Jaquet, CA Mayo, D Osterberg, and MK Marx. 2008. Surveillance, Monitoring and Management of North Atlantic Right Whales in Cape Cod Bay and Adjacent Waters – 2008. October 2008; 186pp, retrieved from <http://www.coastalstudies.org/what-we-do/right-whales/rwreports.htm>

Leeney RH, K Stamieszkin, CA Mayo, and MK Marx. 2009. Surveillance, Monitoring and Management of North Atlantic Right Whales in Cape Cod Bay and Adjacent Waters – 2009. November 2009; 324pp, retrieved from <http://www.coastalstudies.org/what-we-do/right-whales/rwreports.htm>

-
- NMFS (National Marine Fisheries Service). 1991. Recovery Plan for the Humpback Whale (*Megaptera novaeangliae*). Report prepared by the Humpback Whale Recovery Team for the National Marine Fisheries Service, Silver Springs, MD. 105 pp.
- NMFS. 2012. U.S. Atlantic and Gulf of Mexico Marine Mammal Stock Assessments – 2011. NOAA Technical Memorandum NMFS-NE-221. 330 pp. Retrieved from <http://www.nmfs.noaa.gov/pr/sars/region.htm>.
- NMFS. 2013a. Website – “Marine Mammals – Office of Protected Resources” at <http://www.nmfs.noaa.gov/pr/species/mammals/>. Last accessed May 17, 2013.
- NMFS. 2013b. U.S. Atlantic and Gulf of Mexico Marine Mammal Stock Assessments – 2012, volume 1. 425 pp. Retrieved from <http://www.nmfs.noaa.gov/pr/sars/region.htm>.
- MWRA. 2010. Ambient monitoring plan for the Massachusetts Water Resources Authority effluent outfall revision 2. July 2010. Boston: Massachusetts Water Resources Authority. Report 2010-04.
- Payne, PM, Nicolas, J.R., O'Brien, L., and Powers, K.D. 1986. The distribution of the humpback whale, *Megaptera novaeangliae*, on Georges Bank and in the Gulf of Maine in relation to densities of the sand eel, *Ammodytes americanus*. Fish. Bull. U.S.: 271-277.
- Payne PM, DN Wiley, SB Young, S Pittman, PJ Clapham, JW Jossi. 1990. Recent fluctuations in the abundance of baleen whales in the southern Gulf of Maine in relation to changes in selected prey. Fishery Bulletin 88:687-696.
- Pett, S, CJ McKay. 1990. Technical report on the resources and uses of Stellwagen Bank. In: The Resources and Uses of Stellwagen Bank. J.H. Archer (ed.). Urban Harbors Institute, University of Massachusetts, Boston. 66pp.
- Rough V. 1995. Gray seals in Nantucket Sound, Massachusetts, winter and spring, 1994. Final report prepared for the U.S. Marine Mammal Commission. Contract No. T10155615. NTIS No. PB95-191391.
- Schilling, M.R., Seipt, I., Weinrich, M.T., Frohock, S.E., Kuhlberg, A.K., and Clapham, P.J. 1992. Behavior of individually identified sei whales (*Balaenoptera borealis*) during an episodic influx into the southern Gulf of Maine in 1986. Fishery Bulletin 90: 749-755.
- Short, LM, Schaub E. 2006. Summary of marine mammal observations during 2005 surveys. Boston: Massachusetts Water Resources Authority. Report ENQUAD 2005-03. 18 p.
- Short, LM, Michelin D. 2007. Summary of marine mammal observations during 2007 surveys. Boston: Massachusetts Water Resources Authority. Report ENQUAD 2007-01. 19p.
- Stamieszkin K, L Ganley, CA Mayo, RH Leeney, and MK Marx. 2010. Surveillance, Monitoring and Management of North Atlantic Right Whales in Cape Cod Bay and Adjacent Waters – 2010. November 2010; 31pp, retrieved from <http://www.coastalstudies.org/what-we-do/right-whales/rwreports.htm>
- Waring GT, DJ Lalka, PJ Clapham, S Swartz, MC Rossman, TVN Cole, KD Bisack, LJ Hansen. 1999. U.S. Atlantic marine mammal stock assessments-1998. NOAA Technical memorandum NMFS-NE-116.
-

Wenzel, F, DK Matilla, PJ Clapham. 1988. *Balaenoptera musculus* in the Gulf of Maine. Mar. Mamm. Sci. 4(2):172-175.

Weinrich, M., M. Martin, R. Griffiths, J. Bove, and M. Schilling. 1997. A shift in distribution of humpback whales, (*Megaptera novaeangliae*) in response to prey in the southern Gulf of Maine. Fishery Bulletin 95: 826-836.

Weinrich, M.T., C.R. Belt, and D. Morin. 2001. Behavior and ecology of the Atlantic white-sided dolphin (*Lagenorhynchus acutus*) in coastal New England waters. Marine Mammal Science 17: 231-248.

Wu D. 2011. Summary of marine mammal observations during 2010 surveys. Boston: Massachusetts Water Resources Authority. Report 2011-08. 16 p.



Massachusetts Water Resources Authority
Charlestown Navy Yard
100 First Avenue
Boston, MA 02129
(617) 242-6000
<http://www.mwra.state.ma.us>